

## **HIV/STD Statistics**

December 2002

## Relationship of HIV Disease Cases, HIV Cases, and AIDS Cases




From the time a person is infected with human immunodeficiency virus (HIV) until death, he/she has **HIV Disease**. All persons with HIV Disease can be subclassified as <u>either</u> an **AIDS case** (if they are in the later stages of the disease process and have met the case definition for AIDS) <u>or</u> an **HIV case** (if they are in the earlier stages of the disease process and have not met the AIDS case definition).



\* HIV Case vs. AIDS Case


These cases were initially reported as HIV cases, and have subsequently remained HIV cases (i.e., they have not met the case definition for AIDS).



<sup>\*</sup> Based on death certificate data.

<sup>\*\*</sup>Cases are indicated by year of their initial report to the Missouri Department of Health and Senior Services(i.e., by the year in which the first report of the person, whether as an HIV case or an AIDS case, was received by the department.)

<sup>&</sup>quot;These cases were either: 1) initially reported as HIV cases and then later reclassified as AIDS cases because they had subsequently come to meet the AIDS case definition: or 2) initially reported as an AIDS case.




# Reported HIV and AIDS Cases by Gender, Race/Ethnicity, and Age at Diagnosis, Missouri, 1982–2002


	HIV	Cases*	•		AIDS C	ases**		HIV/AIDS	S Cases
Rep	orted 2002	Cum	ulative*	Repoi	rted 2002	Cum	ulative	Cumu	lative
Case	s %	Cases	%	Cases	s %	Cases	%	Cases	%
Gender									
Male250	(79.1%)	3,848	(82.6%)	298	(82.8%)	8,520	(89.9%)	12,368	(87.5%)
Female 66	(20.9%)	809	(17.4%)	62	(17.2%)	958	(10.1%)	1,767	(12.5%)
Race/Ethnicity									
White	(49.4%)	2,504	(53.8%)	185	(51.4%)	6,098	(64.3%)	8,602	(60.9%)
Black	(43.0%)	1,981		160	(44.4%)	3,125	(33.0%)	5,106	(36.1%)
Hispanic 14		114	(2.4%)	14	(3.9%)	197	(2.1%)	311	(2.2%)
Asian/Pacific Islander		16		1		26		42	(0.3%)
American Indian2	` /	15	` /	0		32	,	47	(0.3%)
Unknown6	(1.9%)	27	(0.6%)	0	(0.0%)	0	(0.0%)	27	(0.2%)
Race/Ethnicity and Gender									
White Male131		2,193		168	(46.7%)	5,701		7,894	(55.8%)
Black Male96		1,507		119		2,587		4,094	(29.0%)
Hispanic Male		101		11		181		282	(2.0%)
Asian/Pacific Islander Male		12		0		22		34	(0.2%)
American Indian Male		14		0		29		43	(0.3%)
Unknown Male		21	` ′	0	, ,	0	` ′	21	(0.1%)
White Female		311		17		397	( ,	708	(5.0%)
Black Female		474		41	( ' ' ' ' '	538		1,012	(7.2%)
Hispanic Female 1 Asian/Pacific Islander Female (		13		3	(	16		29	(0.2%)
American Indian Female		4	` /	1 0				8 4	(0.1%) (0.0%)
Unknown Female	(/	6	` /	0		0	,	6	(0.0%)
	(0.070)	0	(0.170)	0	(0.070)	0	(0.070)	0	(0.070)
Age at Diagnosis <sup>‡</sup>									
<13		45		1	(	58	(	103	(0.7%)
13-19	( ,	214		5		101		315	(2.2%)
20-24	(,	728		15	, ,	559	. ,	1,287	(9.1%)
30-34	(14.9%) (24.4%)	1,008		35		1,521		2,527 3,354	(17.9%) (23.7%)
35-39		749	( ' ' ' ' '	87	,	1,982		2,731	(19.3%)
40-44		454		74	,	1,323		1,777	(12.6%)
45-49	()	230		34		760		990	(7.0%)
50+ 16	` /	203	` '	45	` /	848	,	1,051	(7.4%)
Missouri Total 316	(100.0%)	4,657	(100.0%)	360	(100.0%)	9,478	(100.0%)	14,135	(100.0%)


<sup>\*</sup>HIV Cases-Persons with HIV infection who have not developed one of the specific diseases or conditions which would cause them to meet the case definition for AIDS.

\*\*AIDS Cases-Persons with HIV infection who have developed one or more of the specific diseases or conditions which cause them to meet the AIDS case definition.

<sup>&</sup>lt;sup>‡</sup>For HIV Cases, Age at Diagnosis is the age at which the individual was first diagnosed with HIV infection. For AIDS Cases, Age at Diagnosis is the age at which the individual was first diagnosed with AIDS.








Reported HIV and AIDS Cases by Exposure Category, Missouri, 1982–2002										
	HIV Cases*				AIDS Cases**			HIV/AIDS	HIV/AIDS Cases	
	Reported 2002 Cumulative		Repor	Reported 2002 Cum		nulative Cumu		ative		
	Cases	%	Cases	%	Cases	%	Cases	%	Cases	%
Exposure Category <sup>1</sup>										
MSM	120	(38.0%)	2,654	(57.0%)	169	(46.9%)	6,508	(68.7%)	9,162	(64.8%)
MSM/IDU		(3.2%)	273	`	13				1,086	(7.7%)
IDU	10	(3.2%)	393	(8.4%)	28	(7.8%)	709	(7.5%)	1,102	(7.8%)
Heterosexual Contact	35	(11.1%)	748	(16.1%)	47	(13.1%)	837	(8.8%)	1,585	(11.2%)
Adult Hemophiliac	0	(0.0%)	30	(0.6%)	3	(0.8%)	148	(1.6%)	178	(1.3%)
Adult Transfusion	0	(0.0%)	14	(0.3%)	2	(0.6%)	104	(1.1%)	118	(0.8%)
Other/Unknown Adult	140	(44.3%)	500	(10.7%)	94	(26.1%)	287	(3.0%)	787	(5.6%)
Perinatal Transmission	0	(0.0%)	38	(0.8%)	2	(0.6%)	48	(0.5%)	86	(0.6%)
Other/Unknown Pediatric	1	(0.3%)	7	(0.2%)	2	(0.6%)	24	(0.3%)	31	(0.2%)
Missouri Total	. 316 (	(100.0%)	4,657	(100.0%) .	360 (	100.0%)	9,478	(100.0%)	14,135 (	(100.0%)


\*HIV Cases-Persons with HIV infection who have not developed one of the specific diseases or conditions which would cause them to meet the case definition for AIDS.

\*\*AIDS Cases-Persons with HIV infection who have developed one or more of the specific diseases or conditions which cause them to meet the AIDS case definition.

## Reported Adult/Adolescent HIV Cases by Exposure Category<sup>1</sup>, Missouri, 1985-2002



### Reported Adult/Adolescent AIDS Cases by Exposure Category<sup>1</sup>, Missouri, 1982-2002



<sup>1</sup>MSM=men who have sex with men; MSM/IDU=men who have sex with men and inject drugs; IDU=injecting drug users; HETSX=heterosexual contact; Oth/Unk=Other/Unknown

Reported HIV and AIDS Cases and Rates by Area of Residence at Time of Diagnosis,

### Missouri, 1982-2002 **HIV Cases\*** AIDS Cases\*\* Reported 2002 Cumulative Reported 2002 Cumulative Geographic Area Cases Rate\*\*\* Cases % Rate\*\*\* Cases Location St. Louis City† (21.5%) ..... 19.5 1,340 (28.8%)114 (31.7%) ..... 32.7 2,693 (28.4%)St. Louis County<sup>†</sup> (17.1%) ...... 5.3 (13.6%) ..... 4.8 629 (13.5%)(15.5%)54 49 1,466 1,155 76 (24.1%) ..... 17.2 (24.8%)80 (22.2%) ..... 18.1 (27.6%)Kansas City 2,614 Outstate† 86 (27.2%) ...... 2.3 1,190 (25.6%)105 (29.2%) ..... 2.8 2,475 (26.1%)Missouri Correctional Facilities<sup>††</sup> 32 (10.1%) ..... --343 (7.4%)12 (3.3%) ..... --230 (2.4%)Missouri Total 316 (100.0%) ..... 5.6 4,657 (100.0%) 360 (100.0%) ..... 6.4 9,478 (100.0%)

<sup>\*\*</sup>MSM=men who have sex with men; MSM/IDU=men who have sex with men and inject drugs; IDU=injecting drug users


<sup>\*</sup>HIV Cases-Persons with HIV infection who have not developed one of the specific diseases or conditions which would cause them to meet the case definition for AIDS.

<sup>\*\*</sup>AIDS Cases-Persons with HIV infection who have developed one or more of the specific diseases or conditions which cause them to meet the AIDS case definition.

<sup>\*\*\*</sup>Per 100,000 population, based on 2000 population estimates.

<sup>†</sup>Does not include persons living in correctional facilities at the time of diagnosis. These persons are included in the "Missouri Correctional Facilities" category.

 $<sup>^{\</sup>dagger\dagger}\mbox{Includes}$  state, county and local correctional facilities.



Reported HIV Cases by Race/Ethnicity and Area of Residence at Time of Diagnosis, Missouri, 2002						
	Total		White	Black	Hispanic	
Georgraphic Area	2002 Cases	Rate*	2002 Cases Rate*	2002 Cases Rate*	2002 Cases Rate*	
St. Louis City†	68	19.5	23 15.1	37 20.8	3 42.7	
St. Louis County <sup>†</sup>	54	5.3	273.5	24 12.4	2 13.7	
Kansas City†			32 11.9	36 26.1	6 19.6	
Outstate†	86	2.3	611.7	20 16.7	3 4.5	
Missouri Correctional Facilities <sup>††</sup>	32		13	19	0	
Missouri Total* * Per 100 000 population, based on 2000 po			156 3.3	136 21.6	14 11.8	


<sup>\*</sup> Per 100,000 population, based on 2000 population estimates

Currently Living HIV-Diagnosed Persons (HIV and AIDS Cases) Who Were Residents of Missouri at the Time of Diagnosis, and Who Were Reported Through 2002, by Gender and Race/Ethnicity

Living HIV-


	Diagnosed Persons	%
Gender		
Male	7,563 84	.8%
Female	1,355 15	.2%
Race/Ethnicity		
White	5,005 56	.1%
Black	3,617 40	.6%
Hispanic	2112	.4%
Asian/Pacific Islander		
American Indian		
Unknown	27 0	.3%
Race/Ethnicity and Gend	er	
White Male		.4%
Black Male		
Hispanic Male		
Asian/Pacific Islander Male	20 0	.2%
American Indian Male	29 0	.3%
Unknown Male	21 0	.2%
White Female	513 5	.8%
Black Female		
Hispanic Female	25 0	.3%
Asian/Pacific Islander Female	8 0	.1%
American Indian Female		
Unknown Female	6 0	.1%
Total Living HIV-Diagnosed I	Persons8,918 100	.0%

# Currently Living HIV-Diagnosed Persons (HIV and AIDS Cases), Reported Through 2002, by Missouri County of Residence<sup>†</sup> at Time of Diagnosis




Does not include persons living in correctional facilities at the time of diagnosis. These persons are included in the "Missouri Correctional Facilities" category.


<sup>††</sup>Includes state, county and local correctional facilities.




## Reported Gonorrhea Cases and Rates by Geographic Area, Missouri, 2002

Cases	%	Rate*
Missouri		
Whites 1,339	15.0%	28.2
Blacks 6,289	70.3%	999.2
Other/Unknown 1,324	14.8%	
Total Cases 8,952	100.0%	160.0
St. Louis City		
Whites 107	3.9%	70.1
Blacks 2,260	82.6%	1,267.8
Other/Unknown 370	13.5%	
Total Cases 2,737	100.0%	786.1
St. Louis County		
St. Louis County Whites135	7.5%	17.3
Blacks 1.356	75.6%	701.5
Other/Unknown 302	16.8%	701.3
Total Cases 1,793	100.0%	176.4
10tai Cases 1,775	100.0 /0	170.4
Kansas City		
Whites	10.0%	93.3
Blacks 1,921	77.3%	1,393.3
Other/Unknown 315	12.8%	
Total Cases 2,486	100.0%	563.0
Outstate		
Whites 847	43.8%	23.9
Blacks 752	38.8%	627.0
Other/Unknown 337	17.4%	
Total Cases 1,936	100.0%	51.1
*Per 100,000 population		










## Reported P&S Syphilis Cases and Rates by Geographic Area, Missouri, 2002


Cases	s %	Rate*
Missouri		
Whites12	35.3%	0.3
Blacks20	58.8%	3.2
Other/Unknown2	5.9%	
Total Cases34	100.0%	0.6
St. Louis City		
Whites4	30.8%	2.6
Blacks9	69.2%	6.0
Other/Unknown 0	0.0%	
Total Cases13	100.0%	3.7
St. Louis County		
Whites1	14.3%	0.1
Blacks5	71.4%	2.6
Other/Unknown1	14.3%	
Total Cases7	100.0%	0.7
Kansas City		
Whites1	14.3%	0.4
Blacks 5	71.4%	3.6
Other/Unknown1	14.3%	
Total Cases7	100.0%	1.6
Outstate		
Whites6	85.7%	0.2
Blacks 1	14.3%	0.8
Other/Unknown 0	0.0%	
Total Cases7	100.0%	0.2
*Per 100,000 population		





1 Congenital Syphilis case was reported in 2002 in St. Louis City





## Reported Chlamydia Cases and Rates by Geographic Area, Missouri, 2002

	Cases	%	Rate*
Missouri			
Whites	5,474	33.8%	115.3
Blacks		45.0%	1,256.7
Other/Unknown	3,427	21.2%	
Total Cases	16,181	100.0%	289.2
St. Louis City	4.50		
Whites		5.6%	117.2
Blacks		72.2%	1,296.9
Other/Unknown		22.2%	
Total Cases	3,202	100.0%	919.6
St. Louis County	/		
Whites		12.2%	46.8
Blacks	2,012	67.1%	1,040.8
Other/Unknown	623	20.8%	
Total Cases	3,000	100.0%	295.2
Kansas City			
Whites	514	17.5%	191.8
Blacks		61.5%	
Other/Unknown		21.0%	1,312.0
Total Cases			666.3
Total Cases	4,944	100.0 /0	000.3
Outstate			
Whites	4,416	62.8%	124.5
Blacks	1,147	16.3%	956.3
Other/Unknown	1,474	20.9%	
Total Cases	7,037	100.0%	185.7
*Per 100,000 population			

